



United States
Department of
Agriculture


NRCS

Natural
Resources
Conservation
Service

A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for **Rogers County, Oklahoma**

Rogers, County FY19 BUILD Rail Project



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<https://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	8
Soil Map.....	9
Legend.....	10
Map Unit Legend.....	11
Map Unit Descriptions.....	11
Rogers County, Oklahoma.....	14
Bc—Bates-Collinsville complex, 3 to 8 percent slopes.....	14
Br—Eram-Verdigris complex, 0 to 12 percent slopes.....	16
ChB—Choteau silt loam, 1 to 3 percent slopes.....	18
DbC—Dennis-Bates complex, 3 to 5 percent slopes.....	19
DnB—Dennis silt loam, 1 to 3 percent slopes.....	22
DnC—Dennis silt loam, 3 to 5 percent slopes.....	24
PaA—Parsons silt loam, 0 to 1 percent slopes.....	25
TaA—Taloka silt loam, 0 to 1 percent slopes.....	27
Ve—Verdigris clay loam, 0 to 1 percent slopes, occasionally flooded.....	29
Vf—Verdigris silty clay loam, 0 to 2 percent slopes, frequently flooded.....	30
Soil Information for All Uses	32
Suitabilities and Limitations for Use.....	32
Land Classifications.....	32
Farmland Classification (Industrial Lead Enhancement Project - Rogers County).....	32
Soil Reports.....	38
Land Classifications.....	38
Prime and other Important Farmlands (Industrial Lead Enhancement Project - Rogers County).....	38
References	41

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

Custom Soil Resource Report

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and


Custom Soil Resource Report

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.


Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map




Map Scale: 1:25,800 if printed on A landscape (11" x 8.5") sheet.




MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)




















Soils







 Soil Map Unit Polygons

 Soil Map Unit Lines


 Soil Map Unit Points

Special Point Features






-  Blowout
-  Borrow Pit
-  Clay Spot
-  Closed Depression
-  Gravel Pit
-  Gravelly Spot
-  Landfill
-  Lava Flow
-  Marsh or swamp
-  Mine or Quarry
-  Miscellaneous Water
-  Perennial Water
-  Rock Outcrop
-  Saline Spot
-  Sandy Spot
-  Severely Eroded Spot
-  Sinkhole
-  Slide or Slip
-  Sodic Spot

-  Spoil Area
-  Stony Spot
-  Very Stony Spot
-  Wet Spot
-  Other
-  Special Line Features


Water Features

 Streams and Canals

Transportation

-  Rails
-  Interstate Highways
-  US Routes
-  Major Roads
-  Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Rogers County, Oklahoma
 Survey Area Data: Version 13, Sep 5, 2018

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Feb 14, 2015—Nov 25, 2017

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
Bc	Bates-Collinsville complex, 3 to 8 percent slopes	4.1	2.4%
Br	Eram-Verdigris complex, 0 to 12 percent slopes	15.4	9.1%
ChB	Choteau silt loam, 1 to 3 percent slopes	83.4	49.2%
DbC	Dennis-Bates complex, 3 to 5 percent slopes	17.8	10.5%
DnB	Dennis silt loam, 1 to 3 percent slopes	16.2	9.5%
DnC	Dennis silt loam, 3 to 5 percent slopes	6.3	3.7%
PaA	Parsons silt loam, 0 to 1 percent slopes	3.3	2.0%
TaA	Taloka silt loam, 0 to 1 percent slopes	16.3	9.6%
Ve	Verdigris clay loam, 0 to 1 percent slopes, occasionally flooded	1.8	1.0%
Vf	Verdigris silty clay loam, 0 to 2 percent slopes, frequently flooded	5.0	3.0%
Totals for Area of Interest		169.6	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a

Custom Soil Resource Report

particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Custom Soil Resource Report

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Rogers County, Oklahoma

Bc—Bates-Collinsville complex, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2xvgn
Elevation: 510 to 950 feet
Mean annual precipitation: 41 to 45 inches
Mean annual air temperature: 57 to 61 degrees F
Frost-free period: 170 to 235 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Bates and similar soils: 70 percent
Collinsville and similar soils: 20 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Bates

Setting

Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Residuum weathered from sandstone and shale

Typical profile

A - 0 to 13 inches: loam
BA - 13 to 19 inches: loam
Bt - 19 to 31 inches: clay loam
Cr - 31 to 42 inches: bedrock

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 26 to 38 inches to paralithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: Moderate (about 6.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4e
Hydrologic Soil Group: C
Ecological site: Loamy Upland (R112XY103KS)
Hydric soil rating: No

Description of Collinsville

Setting

Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Residuum weathered from sandstone

Typical profile

A - 0 to 8 inches: loam
C - 8 to 12 inches: gravelly fine sandy loam
R - 12 to 22 inches: bedrock

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 11 to 20 inches to lithic bedrock
Natural drainage class: Somewhat excessively drained
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately high (0.00 to 0.20 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: Very low (about 1.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4s
Hydrologic Soil Group: D
Ecological site: Shallow Sandstone Upland (R112XY105OK)
Hydric soil rating: No

Minor Components

Dennis

Percent of map unit: 5 percent
Landform: Hillslopes
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Base slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: Loamy Upland (R112XY103KS)
Hydric soil rating: No

Parsons

Percent of map unit: 5 percent
Landform: Divides
Landform position (two-dimensional): Summit
Landform position (three-dimensional): Interfluve
Down-slope shape: Concave
Across-slope shape: Concave
Ecological site: Claypan Upland (R112XY101KS)
Hydric soil rating: No

Br—Eram-Verdigris complex, 0 to 12 percent slopes

Map Unit Setting

National map unit symbol: 2tgtn
Elevation: 520 to 1,020 feet
Mean annual precipitation: 39 to 47 inches
Mean annual air temperature: 57 to 63 degrees F
Frost-free period: 200 to 220 days
Farmland classification: Not prime farmland

Map Unit Composition

Eram and similar soils: 55 percent
Verdigris and similar soils: 30 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Eram

Setting

Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Clayey residuum weathered from sandstone and shale

Typical profile

A - 0 to 12 inches: silt loam
Bt - 12 to 24 inches: silty clay
BC - 24 to 36 inches: silty clay
Cr - 36 to 46 inches: bedrock

Properties and qualities

Slope: 0 to 12 percent
Depth to restrictive feature: 20 to 40 inches to paralithic bedrock
Natural drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately high (0.00 to 0.20 in/hr)
Depth to water table: About 16 to 30 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: Moderate (about 6.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 6e
Hydrologic Soil Group: D

Custom Soil Resource Report

Ecological site: Loamy prairie PE 62-80 (R112XY056OK)
Hydric soil rating: No

Description of Verdigris

Setting

Landform: Drainageways
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Silty alluvium

Typical profile

A - 0 to 22 inches: silt loam
AC - 22 to 42 inches: silt loam
C - 42 to 79 inches: silt loam

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: Frequent
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: High (about 12.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 5w
Hydrologic Soil Group: B
Ecological site: Loamy Lowland (Draft) (PE 35-42) (R112XY013KS)
Hydric soil rating: No

Minor Components

Bates

Percent of map unit: 5 percent
Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: Loamy prairie (Northeast) PE 62-80 (R112XY059OK)
Hydric soil rating: No

Collinsville

Percent of map unit: 5 percent
Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: Shallow Sandstone (Draft) (PE 35-42) (R112XY030KS)

Custom Soil Resource Report

Hydric soil rating: No

Taloka

Percent of map unit: 3 percent

Landform: Paleoterraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Linear

Ecological site: Loamy prairie (Northeast) PE 62-80 (R112XY059OK)

Hydric soil rating: No

Summit

Percent of map unit: 2 percent

Landform: Interfluves

Landform position (two-dimensional): Backslope, footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave, convex

Across-slope shape: Concave

Ecological site: Loamy prairie (Northeast) PE 62-80 (R112XY059OK)

Hydric soil rating: No

ChB—Choteau silt loam, 1 to 3 percent slopes

Map Unit Setting

National map unit symbol: 2tgtk

Elevation: 490 to 1,120 feet

Mean annual precipitation: 37 to 48 inches

Mean annual air temperature: 57 to 61 degrees F

Frost-free period: 190 to 220 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Choteau and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Choteau

Setting

Landform: Paleoterraces

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Convex

Parent material: Loamy and clayey alluvium and/or colluvium over residuum weathered from shale

Typical profile

A - 0 to 15 inches: silt loam

E - 15 to 24 inches: silt loam

BE - 24 to 31 inches: silty clay loam

Custom Soil Resource Report

Bt1 - 31 to 40 inches: clay

Bt2 - 40 to 65 inches: clay

Properties and qualities

Slope: 1 to 3 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 30 to 36 inches

Frequency of flooding: None

Frequency of ponding: None

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water storage in profile: High (about 10.8 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2e

Hydrologic Soil Group: C

Ecological site: Loamy Upland (R112XY103KS)

Hydric soil rating: No

Minor Components

Bucyrus

Percent of map unit: 5 percent

Landform: Interfluves

Landform position (two-dimensional): Shoulder, summit

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: Loamy Upland (R112XY103KS)

Hydric soil rating: No

Taloka

Percent of map unit: 5 percent

Landform: Paleoterraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Linear

Ecological site: Claypan Upland (R112XY101KS)

Hydric soil rating: No

DbC—Dennis-Bates complex, 3 to 5 percent slopes

Map Unit Setting

National map unit symbol: 2wqf1

Elevation: 540 to 930 feet

Mean annual precipitation: 41 to 45 inches

Mean annual air temperature: 57 to 61 degrees F

Custom Soil Resource Report

Frost-free period: 190 to 220 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Dennis and similar soils: 50 percent

Bates and similar soils: 40 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Dennis

Setting

Landform: Hillslopes

Landform position (two-dimensional): Footslope, backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Silty and clayey residuum weathered from shale

Typical profile

A - 0 to 8 inches: silt loam

BA - 8 to 20 inches: silty clay loam

Bt1 - 20 to 25 inches: silty clay

Bt2 - 25 to 62 inches: silty clay

C - 62 to 79 inches: silty clay loam

Properties and qualities

Slope: 3 to 5 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 12 to 24 inches

Frequency of flooding: None

Frequency of ponding: None

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water storage in profile: High (about 9.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: C/D

Ecological site: Loamy Upland (R112XY103KS)

Hydric soil rating: No

Description of Bates

Setting

Landform: Hillslopes

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Residuum weathered from sandstone and shale

Custom Soil Resource Report

Typical profile

A - 0 to 13 inches: loam
BA - 13 to 19 inches: loam
Bt - 19 to 31 inches: clay loam
Cr - 31 to 42 inches: bedrock

Properties and qualities

Slope: 3 to 5 percent
Depth to restrictive feature: 26 to 38 inches to paralithic bedrock
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: Moderate (about 6.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: C
Ecological site: Loamy Upland (R112XY103KS)
Hydric soil rating: No

Minor Components

Collinsville

Percent of map unit: 5 percent
Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: Shallow Sandstone Upland (R112XY105OK)
Hydric soil rating: No

Parsons

Percent of map unit: 5 percent
Landform: Paleoterraces
Landform position (three-dimensional): Tread
Down-slope shape: Concave
Across-slope shape: Concave
Ecological site: Claypan Upland (R112XY101KS)
Hydric soil rating: No

DnB—Dennis silt loam, 1 to 3 percent slopes

Map Unit Setting

National map unit symbol: 2tgsq
Elevation: 460 to 1,260 feet
Mean annual precipitation: 37 to 45 inches
Mean annual air temperature: 55 to 61 degrees F
Frost-free period: 150 to 255 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Dennis and similar soils: 82 percent
Minor components: 18 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Dennis

Setting

Landform: Interfluves
Landform position (two-dimensional): Footslope, summit
Landform position (three-dimensional): Base slope, interfluve
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Silty and clayey residuum weathered from shale

Typical profile

A - 0 to 11 inches: silt loam
BA - 11 to 17 inches: silty clay loam
Bt1 - 17 to 22 inches: silty clay
Bt2 - 22 to 68 inches: silty clay
C - 68 to 79 inches: silty clay loam

Properties and qualities

Slope: 1 to 3 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: About 12 to 24 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: Moderate (about 8.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 2e
Hydrologic Soil Group: C/D

Custom Soil Resource Report

Ecological site: Loamy Upland (R112XY103KS)

Hydric soil rating: No

Minor Components

Bates

Percent of map unit: 5 percent

Landform: Interfluves

Landform position (two-dimensional): Summit, shoulder

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: Loamy Upland (R112XY103KS)

Hydric soil rating: No

Parsons

Percent of map unit: 5 percent

Landform: Divides

Landform position (two-dimensional): Summit

Landform position (three-dimensional): Interfluve

Down-slope shape: Concave

Across-slope shape: Concave

Ecological site: Claypan Upland (R112XY101KS)

Hydric soil rating: No

Eram

Percent of map unit: 5 percent

Landform: Interfluves

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: Loamy Upland (R112XY103KS)

Hydric soil rating: No

Kenoma

Percent of map unit: 2 percent

Landform: Interfluves

Landform position (two-dimensional): Summit

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: Clayey Upland (R112XY102KS)

Hydric soil rating: No

Pharoah

Percent of map unit: 1 percent

Landform: Paleoterraces

Landform position (three-dimensional): Tread

Down-slope shape: Concave

Across-slope shape: Linear

Ecological site: Claypan Upland (R112XY101KS)

Hydric soil rating: No

DnC—Dennis silt loam, 3 to 5 percent slopes

Map Unit Setting

National map unit symbol: 2thf0
Elevation: 600 to 1,000 feet
Mean annual precipitation: 41 to 45 inches
Mean annual air temperature: 55 to 61 degrees F
Frost-free period: 150 to 255 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Dennis and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Dennis

Setting

Landform: Hillslopes
Landform position (two-dimensional): Backslope, shoulder
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Silty and clayey residuum weathered from shale

Typical profile

A - 0 to 11 inches: silt loam
BA - 11 to 17 inches: silty clay loam
Bt1 - 17 to 22 inches: silty clay
Bt2 - 22 to 68 inches: silty clay
C - 68 to 79 inches: silty clay loam

Properties and qualities

Slope: 3 to 5 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: About 12 to 24 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: High (about 10.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: C/D
Ecological site: Loamy Upland (R112XY103KS)

Custom Soil Resource Report

Hydric soil rating: No

Minor Components

Bates

Percent of map unit: 5 percent

Landform: Interfluves

Landform position (two-dimensional): Shoulder, backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: Loamy Upland (R112XY103KS)

Hydric soil rating: No

Collinsville

Percent of map unit: 4 percent

Landform: Hillslopes

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: Shallow Sandstone Upland (R112XY105OK)

Hydric soil rating: No

Eram

Percent of map unit: 3 percent

Landform: Interfluves

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: Clayey Upland (R112XY102KS)

Hydric soil rating: No

Parsons

Percent of map unit: 3 percent

Landform: Interfluves

Landform position (two-dimensional): Summit

Landform position (three-dimensional): Interfluve

Down-slope shape: Concave

Across-slope shape: Concave

Ecological site: Claypan Upland (R112XY101KS)

Hydric soil rating: No

PaA—Parsons silt loam, 0 to 1 percent slopes

Map Unit Setting

National map unit symbol: 2thdx

Elevation: 510 to 1,340 feet

Mean annual precipitation: 35 to 45 inches

Mean annual air temperature: 55 to 61 degrees F

Custom Soil Resource Report

Frost-free period: 175 to 230 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Parsons and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Parsons

Setting

Landform: Divides

Landform position (two-dimensional): Summit

Landform position (three-dimensional): Interfluve

Down-slope shape: Concave

Across-slope shape: Concave

Parent material: Loess over clayey alluvium and/or clayey residuum weathered from clayey shale

Typical profile

Ap - 0 to 8 inches: silt loam

E - 8 to 14 inches: silt loam

2Btg1 - 14 to 24 inches: silty clay

2Btg2 - 24 to 39 inches: silty clay

2BC - 39 to 59 inches: silty clay loam

2C - 59 to 79 inches: silty clay loam

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: 9 to 17 inches to abrupt textural change

Natural drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: About 6 to 18 inches

Frequency of flooding: None

Frequency of ponding: None

Gypsum, maximum in profile: 6 percent

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water storage in profile: Very low (about 2.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3w

Hydrologic Soil Group: D

Ecological site: Claypan Summit Prairie (R112XY011MO)

Hydric soil rating: No

Minor Components

Dennis

Percent of map unit: 7 percent

Landform: Interfluves

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Convex

Across-slope shape: Convex

Custom Soil Resource Report

Ecological site: Loamy Upland (Draft) (PE 35-42) (R112XY015KS)
Hydric soil rating: No

Pharoah

Percent of map unit: 5 percent
Landform: Paleoterraces
Landform position (three-dimensional): Tread
Down-slope shape: Concave
Across-slope shape: Linear
Ecological site: Claypan prairie PE 62-80 (R112XY010OK)
Hydric soil rating: No

Zaar

Percent of map unit: 3 percent
Landform: Interfluves
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Base slope
Down-slope shape: Convex
Across-slope shape: Convex
Ecological site: Clay Upland (PE 35-42) (R112XY007KS)
Hydric soil rating: No

Aquolls

Percent of map unit: 0 percent
Landform: Divides
Landform position (two-dimensional): Summit
Landform position (three-dimensional): Interfluve
Down-slope shape: Concave
Across-slope shape: Concave
Ecological site: Clay Upland (PE 35-42) (R112XY007KS)
Hydric soil rating: Yes

TaA—Taloka silt loam, 0 to 1 percent slopes

Map Unit Setting

National map unit symbol: 2thf3
Elevation: 500 to 1,200 feet
Mean annual precipitation: 37 to 45 inches
Mean annual air temperature: 54 to 63 degrees F
Frost-free period: 185 to 255 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Taloka and similar soils: 92 percent
Minor components: 8 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Taloka

Setting

Landform: Paleoterraces

Custom Soil Resource Report

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy and clayey alluvium and/or loamy and clayey colluvium over residuum weathered from sandstone and shale

Typical profile

Ap - 0 to 8 inches: silt loam

E - 8 to 20 inches: silt loam

2Btg1 - 20 to 24 inches: silty clay

2Btg2 - 24 to 39 inches: silty clay

2BC - 39 to 59 inches: silty clay loam

2C - 59 to 79 inches: silty clay loam

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: 9 to 24 inches to abrupt textural change

Natural drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately low (0.00 to 0.06 in/hr)

Depth to water table: About 6 to 18 inches

Frequency of flooding: None

Frequency of ponding: None

Gypsum, maximum in profile: 6 percent

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water storage in profile: Low (about 4.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3s

Hydrologic Soil Group: D

Ecological site: Loamy prairie (Northeast) PE 62-80 (R112XY059OK)

Hydric soil rating: No

Minor Components

Dennis

Percent of map unit: 5 percent

Landform: Interfluves

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: Loamy Upland (Draft) (PE 35-42) (R112XY015KS)

Hydric soil rating: No

Bates

Percent of map unit: 3 percent

Landform: Interfluves

Landform position (two-dimensional): Summit, shoulder

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Convex

Ecological site: Sandstone/Shale Upland Prairie (R112XY016MO)

Hydric soil rating: No

Aquolls

Percent of map unit: 0 percent
Landform: Divides
Landform position (two-dimensional): Summit
Landform position (three-dimensional): Interfluve
Down-slope shape: Concave
Across-slope shape: Concave
Ecological site: Clay Upland (PE 35-42) (R112XY007KS)
Hydric soil rating: Yes

Ve—Verdigris clay loam, 0 to 1 percent slopes, occasionally flooded

Map Unit Setting

National map unit symbol: 2tgsn
Elevation: 510 to 890 feet
Mean annual precipitation: 37 to 45 inches
Mean annual air temperature: 57 to 61 degrees F
Frost-free period: 178 to 235 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Verdigris and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Verdigris

Setting

Landform: Flood plains
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Silty alluvium

Typical profile

A - 0 to 20 inches: clay loam
C - 20 to 79 inches: silty clay loam

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.20 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: Occasional
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: High (about 10.8 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 2w
Hydrologic Soil Group: C
Ecological site: Loamy Lowland (Draft) (PE 35-42) (R112XY013KS)
Hydric soil rating: No

Minor Components

Osage, hydric

Percent of map unit: 5 percent
Landform: Flood plains
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Clay Lowland (PE 35-42) (R112XY004KS)
Hydric soil rating: Yes

Cleora

Percent of map unit: 4 percent
Landform: Flood plains
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Quercus macrocarpa-Quercus bicolor/Elymus virginicus-Cinna arundinacea (F112XY052MO)
Hydric soil rating: No

Tallahassee

Percent of map unit: 1 percent
Landform: Flood plains
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Ecological site: Subirrigated PE 62-80 (R112XY095OK)
Hydric soil rating: No

Vf—Verdigris silty clay loam, 0 to 2 percent slopes, frequently flooded

Map Unit Setting

National map unit symbol: 310h
Elevation: 500 to 1,000 feet
Mean annual precipitation: 38 to 47 inches
Mean annual air temperature: 57 to 64 degrees F
Frost-free period: 190 to 230 days
Farmland classification: Not prime farmland

Map Unit Composition

Verdigris and similar soils: 100 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Verdigris

Setting

Landform: Flood plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Silty alluvium

Typical profile

A - 0 to 17 inches: silty clay loam
C - 17 to 80 inches: silty clay loam

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: Frequent
Frequency of ponding: None
Available water storage in profile: High (about 12.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 5w
Hydrologic Soil Group: B
Ecological site: Loamy bottomland PE 62-80 (R112XY050OK)
Forage suitability group: Unnamed (G112XY057OK)
Hydric soil rating: No

Soil Information for All Uses

Suitabilities and Limitations for Use

The Suitabilities and Limitations for Use section includes various soil interpretations displayed as thematic maps with a summary table for the soil map units in the selected area of interest. A single value or rating for each map unit is generated by aggregating the interpretive ratings of individual map unit components. This aggregation process is defined for each interpretation.


Land Classifications

Land Classifications are specified land use and management groupings that are assigned to soil areas because combinations of soil have similar behavior for specified practices. Most are based on soil properties and other factors that directly influence the specific use of the soil. Example classifications include ecological site classification, farmland classification, irrigated and nonirrigated land capability classification, and hydric rating.


Farmland Classification (Industrial Lead Enhancement Project - Rogers County)

Farmland classification identifies map units as prime farmland, farmland of statewide importance, farmland of local importance, or unique farmland. It identifies the location and extent of the soils that are best suited to food, feed, fiber, forage, and oilseed crops. NRCS policy and procedures on prime and unique farmlands are published in the "Federal Register," Vol. 43, No. 21, January 31, 1978.

Custom Soil Resource Report
 Map—Farmland Classification (Industrial Lead Enhancement Project - Rogers County)



Map Scale: 1:25,800 if printed on A landscape (11" x 8.5") sheet.




Map projection: Web Mercator Corner coordinates: WGS84 Edge tics: UTM Zone 15N WGS84

Custom Soil Resource Report









MAP LEGEND








Area of Interest (AOI)






 Area of Interest (AOI)








Soils



Soil Rating Polygons

-  Not prime farmland
-  All areas are prime farmland
-  Prime farmland if drained
-  Prime farmland if protected from flooding or not frequently flooded during the growing season
-  Prime farmland if irrigated
-  Prime farmland if drained and either protected from flooding or not frequently flooded during the growing season
-  Prime farmland if irrigated and drained
-  Prime farmland if irrigated and either protected from flooding or not frequently flooded during the growing season









-  Prime farmland if subsoiled, completely removing the root inhibiting soil layer
-  Prime farmland if irrigated and the product of I (soil erodibility) x C (climate factor) does not exceed 60
-  Prime farmland if irrigated and reclaimed of excess salts and sodium
-  Farmland of statewide importance
-  Farmland of statewide importance, if drained
-  Farmland of statewide importance, if protected from flooding or not frequently flooded during the growing season
-  Farmland of statewide importance, if irrigated

-  Farmland of statewide importance, if drained and either protected from flooding or not frequently flooded during the growing season
-  Farmland of statewide importance, if irrigated and drained
-  Farmland of statewide importance, if irrigated and either protected from flooding or not frequently flooded during the growing season
-  Farmland of statewide importance, if subsoiled, completely removing the root inhibiting soil layer
-  Farmland of statewide importance, if irrigated and the product of I (soil erodibility) x C (climate factor) does not exceed 60







































-  Farmland of statewide importance, if irrigated and reclaimed of excess salts and sodium
-  Farmland of statewide importance, if drained or either protected from flooding or not frequently flooded during the growing season
-  Farmland of statewide importance, if warm enough, and either drained or either protected from flooding or not frequently flooded during the growing season
-  Farmland of statewide importance, if warm enough
-  Farmland of statewide importance, if thawed
-  Farmland of local importance
-  Farmland of local importance, if irrigated

-  Farmland of unique importance
-  Not rated or not available

Soil Rating Lines

-  Not prime farmland
-  All areas are prime farmland
-  Prime farmland if drained
-  Prime farmland if protected from flooding or not frequently flooded during the growing season
-  Prime farmland if irrigated
-  Prime farmland if drained and either protected from flooding or not frequently flooded during the growing season
-  Prime farmland if irrigated and drained
-  Prime farmland if irrigated and either protected from flooding or not frequently flooded during the growing season

Custom Soil Resource Report

	Prime farmland if subsoiled, completely removing the root inhibiting soil layer		Farmland of statewide importance, if drained and either protected from flooding or not frequently flooded during the growing season		Farmland of statewide importance, if irrigated and reclaimed of excess salts and sodium		Farmland of unique importance		Prime farmland if subsoiled, completely removing the root inhibiting soil layer
	Prime farmland if irrigated and the product of I (soil erodibility) x C (climate factor) does not exceed 60		Farmland of statewide importance, if irrigated and drained		Farmland of statewide importance, if drained or either protected from flooding or not frequently flooded during the growing season		Not prime farmland		Prime farmland if irrigated and the product of I (soil erodibility) x C (climate factor) does not exceed 60
	Prime farmland if irrigated and reclaimed of excess salts and sodium		Farmland of statewide importance, if irrigated and either protected from flooding or not frequently flooded during the growing season		Farmland of statewide importance, if warm enough, and either drained or either protected from flooding or not frequently flooded during the growing season		Prime farmland if drained		Prime farmland if irrigated and reclaimed of excess salts and sodium
	Farmland of statewide importance		Farmland of statewide importance, if irrigated and either protected from flooding or not frequently flooded during the growing season		Farmland of statewide importance, if warm enough, and either drained or either protected from flooding or not frequently flooded during the growing season		Prime farmland if protected from flooding or not frequently flooded during the growing season		Farmland of statewide importance
	Farmland of statewide importance, if drained		Farmland of statewide importance, if subsoiled, completely removing the root inhibiting soil layer		Farmland of statewide importance, if subsoiled, completely removing the root inhibiting soil layer		Prime farmland if irrigated		Farmland of statewide importance, if drained
	Farmland of statewide importance, if protected from flooding or not frequently flooded during the growing season		Farmland of statewide importance, if irrigated and the product of I (soil erodibility) x C (climate factor) does not exceed 60		Farmland of statewide importance, if warm enough		Prime farmland if drained and either protected from flooding or not frequently flooded during the growing season		Farmland of statewide importance, if protected from flooding or not frequently flooded during the growing season
	Farmland of statewide importance, if irrigated				Farmland of statewide importance, if thawed		Prime farmland if irrigated and drained		Farmland of statewide importance, if irrigated
					Farmland of local importance		Prime farmland if irrigated and either protected from flooding or not frequently flooded during the growing season		
					Farmland of local importance, if irrigated				

Custom Soil Resource Report

<ul style="list-style-type: none"> Farmland of statewide importance, if drained and either protected from flooding or not frequently flooded during the growing season Farmland of statewide importance, if irrigated and drained Farmland of statewide importance, if irrigated and either protected from flooding or not frequently flooded during the growing season Farmland of statewide importance, if subsoiled, completely removing the root inhibiting soil layer Farmland of statewide importance, if irrigated and the product of I (soil erodibility) x C (climate factor) does not exceed 60 	<ul style="list-style-type: none"> Farmland of statewide importance, if irrigated and reclaimed of excess salts and sodium Farmland of statewide importance, if drained or either protected from flooding or not frequently flooded during the growing season Farmland of statewide importance, if warm enough, and either drained or either protected from flooding or not frequently flooded during the growing season Farmland of statewide importance, if warm enough Farmland of statewide importance, if thawed Farmland of local importance Farmland of local importance, if irrigated 	<ul style="list-style-type: none"> Farmland of unique importance Not rated or not available <p>Water Features</p> <ul style="list-style-type: none"> Streams and Canals <p>Transportation</p> <ul style="list-style-type: none"> Rails Interstate Highways US Routes Major Roads Local Roads <p>Background</p> <ul style="list-style-type: none"> Aerial Photography 	<p>The soil surveys that comprise your AOI were mapped at 1:24,000.</p> <p>Please rely on the bar scale on each map sheet for map measurements.</p> <p>Source of Map: Natural Resources Conservation Service Web Soil Survey URL: Coordinate System: Web Mercator (EPSG:3857)</p> <p>Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.</p> <p>This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.</p> <p>Soil Survey Area: Rogers County, Oklahoma Survey Area Data: Version 13, Sep 5, 2018</p> <p>Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.</p> <p>Date(s) aerial images were photographed: Feb 14, 2015—Nov 25, 2017</p> <p>The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.</p>
---	---	---	---

Table—Farmland Classification (Industrial Lead Enhancement Project - Rogers County)

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
Bc	Bates-Collinsville complex, 3 to 8 percent slopes	All areas are prime farmland	4.1	2.4%
Br	Eram-Verdigris complex, 0 to 12 percent slopes	Not prime farmland	15.4	9.1%
ChB	Choteau silt loam, 1 to 3 percent slopes	All areas are prime farmland	83.4	49.2%
DbC	Dennis-Bates complex, 3 to 5 percent slopes	All areas are prime farmland	17.8	10.5%
DnB	Dennis silt loam, 1 to 3 percent slopes	All areas are prime farmland	16.2	9.5%
DnC	Dennis silt loam, 3 to 5 percent slopes	All areas are prime farmland	6.3	3.7%
PaA	Parsons silt loam, 0 to 1 percent slopes	All areas are prime farmland	3.3	2.0%
TaA	Taloka silt loam, 0 to 1 percent slopes	All areas are prime farmland	16.3	9.6%
Ve	Verdigris clay loam, 0 to 1 percent slopes, occasionally flooded	All areas are prime farmland	1.8	1.0%
Vf	Verdigris silty clay loam, 0 to 2 percent slopes, frequently flooded	Not prime farmland	5.0	3.0%
Totals for Area of Interest			169.6	100.0%

Rating Options—Farmland Classification (Industrial Lead Enhancement Project - Rogers County)

Aggregation Method: No Aggregation Necessary

Tie-break Rule: Lower

Soil Reports

The Soil Reports section includes various formatted tabular and narrative reports (tables) containing data for each selected soil map unit and each component of each unit. No aggregation of data has occurred as is done in reports in the Soil Properties and Qualities and Suitabilities and Limitations sections.

The reports contain soil interpretive information as well as basic soil properties and qualities. A description of each report (table) is included.

Land Classifications

This folder contains a collection of tabular reports that present a variety of soil groupings. The reports (tables) include all selected map units and components for each map unit. Land classifications are specified land use and management groupings that are assigned to soil areas because combinations of soil have similar behavior for specified practices. Most are based on soil properties and other factors that directly influence the specific use of the soil. Example classifications include ecological site classification, farmland classification, irrigated and nonirrigated land capability classification, and hydric rating.

Prime and other Important Farmlands (Industrial Lead Enhancement Project - Rogers County)

This table lists the map units in the survey area that are considered important farmlands. Important farmlands consist of prime farmland, unique farmland, and farmland of statewide or local importance. This list does not constitute a recommendation for a particular land use.

In an effort to identify the extent and location of important farmlands, the Natural Resources Conservation Service, in cooperation with other interested Federal, State, and local government organizations, has inventoried land that can be used for the production of the Nation's food supply.

Prime farmland is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil quality, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and

Custom Soil Resource Report

growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. The water supply is dependable and of adequate quality. Prime farmland is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

For some of the soils identified in the table as prime farmland, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures.

A recent trend in land use in some areas has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

Unique farmland is land other than prime farmland that is used for the production of specific high-value food and fiber crops, such as citrus, tree nuts, olives, cranberries, and other fruits and vegetables. It has the special combination of soil quality, growing season, moisture supply, temperature, humidity, air drainage, elevation, and aspect needed for the soil to economically produce sustainable high yields of these crops when properly managed. The water supply is dependable and of adequate quality. Nearness to markets is an additional consideration. Unique farmland is not based on national criteria. It commonly is in areas where there is a special microclimate, such as the wine country in California.

In some areas, land that does not meet the criteria for prime or unique farmland is considered to be *farmland of statewide importance* for the production of food, feed, fiber, forage, and oilseed crops. The criteria for defining and delineating farmland of statewide importance are determined by the appropriate State agencies. Generally, this land includes areas of soils that nearly meet the requirements for prime farmland and that economically produce high yields of crops when treated and managed according to acceptable farming methods. Some areas may produce as high a yield as prime farmland if conditions are favorable. Farmland of statewide importance may include tracts of land that have been designated for agriculture by State law.

In some areas that are not identified as having national or statewide importance, land is considered to be *farmland of local importance* for the production of food, feed, fiber, forage, and oilseed crops. This farmland is identified by the appropriate local agencies. Farmland of local importance may include tracts of land that have been designated for agriculture by local ordinance.

Report—Prime and other Important Farmlands (Industrial Lead Enhancement Project - Rogers County)

Prime and other Important Farmlands—Rogers County, Oklahoma		
Map Symbol	Map Unit Name	Farmland Classification
Bc	Bates-Collinsville complex, 3 to 8 percent slopes	All areas are prime farmland
Br	Eram-Verdigris complex, 0 to 12 percent slopes	Not prime farmland
ChB	Choteau silt loam, 1 to 3 percent slopes	All areas are prime farmland

Custom Soil Resource Report

Prime and other Important Farmlands–Rogers County, Oklahoma		
Map Symbol	Map Unit Name	Farmland Classification
DbC	Dennis-Bates complex, 3 to 5 percent slopes	All areas are prime farmland
DnB	Dennis silt loam, 1 to 3 percent slopes	All areas are prime farmland
DnC	Dennis silt loam, 3 to 5 percent slopes	All areas are prime farmland
PaA	Parsons silt loam, 0 to 1 percent slopes	All areas are prime farmland
TaA	Taloka silt loam, 0 to 1 percent slopes	All areas are prime farmland
Ve	Verdigris clay loam, 0 to 1 percent slopes, occasionally flooded	All areas are prime farmland
Vf	Verdigris silty clay loam, 0 to 2 percent slopes, frequently flooded	Not prime farmland

References

- American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.
- American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.
- Federal Register. July 13, 1994. Changes in hydric soils of the United States.
- Federal Register. September 18, 2002. Hydric soils of the United States.
- Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.
- National Research Council. 1995. Wetlands: Characteristics and boundaries.
- Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_054262
- Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577
- Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580
- Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.
- United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.
- United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053374
- United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. <http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084>

Custom Soil Resource Report

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf